

Sedgwick County Medical Society

Core Measures Update:

1Q08 - 2Q08

	Indicator	Galichia Medical Center		Kansas Heart Hospital		Kansas Spine Hospital		Via Christi		Wesley Medical Center		Kansas Medical Center		Ntl 90th %
		2Q08	3Q08	2Q08	3Q08	2Q08	3Q08	2Q08	3Q08	2Q08	3Q08	2Q08	3Q08	3Q07 - 2Q08
AMI-1	Aspirin at arrival	100% (10)	100% (12)	100% (6)	100% (9)			95% (59)	99% (68)	100% (60)	100% (66)	100% (6)	100% (9)	100%
AMI-2	Aspirin prescribed at discharge	90% (29)	100% (27)	100% (45)	98% (46)			98% (80)	95% (76)	100% (88)	97% (92)	97% (29)	100% (22)	100%
AMI-3	ACEI or ARB for LVSD	50% (4)	100% (6)	100% (3)	90% (10)			94% (17)	80% (10)	100% (10)	100% (17)	100% (4)	100% (3)	100%
AMI-4	Adult smoking cessation advice/counseling	91% (11)	100% (12)	93% (14)	100% (21)			100%	100%	100% (34)	100% (40)	100% (10)	100% (8)	100%
AMI-5	Beta blocker prescribed at discharge	85% (11)	92% (24)	97% (39)	100% (38)			98% (81)	99% (73)	100% (59)	95% (64)	94% (32)	96% (23)	100%
AMI-6	Beta blocker at arrival (Measure To Be Retired 1 April 2009)	47% (15)	86% (14)	80% (5)	100% (7)			96% (53)	97% (58)	97% (34)	96% (44)	100% (6)	100% (8)	NA
AMI-7A	Fibrinolytic therapy received within 30 minutes of hospital arrival													100%
AMI-8A	PCI received within 90 minutes of hospital arrival	50% (4)	43% (7)					50% (10)	50% (16)	91% (11)	100% (19)	100% (2)	100% (3)	94%
PN-1	Oxygenation assessment (Measure Retired Effective 1 January 2009)	100% (27)	100% (19)					100%	100%	100% (74)	100% (45)	100% (10)	100% (2)	100%
PN-2	Pneumococcal vaccination	66% (29)	85% (13)					55% (44)	45% (38)	97% (67)	97% (38)	69% (13)	100% (2)	98%
PN-3A	Blood cultures performed within 24 hours prior to or 24 hours after hospital arrival for patients who were transferred or admitted to the ICU		100%					100%	100%	96% (26)	100% (15)	0 NA	0 NA	NA
PN-3B	Blood cultures performed in the ED prior to initial antibiotic received in hospital	83% (18)	92% (12)					90% (39)	88% (34)	98% (42)	100% (23)	100% (7)	0 NA	100%
PN-4	Adult smoking cessation advice/counseling	88% (8)	100% (7)					100%	94% (16)	100% (37)	100% (19)	100% (1)	0 NA	100%
PN-5B	Initial antibiotic received within 6 hours of hospital arrival	78% (18)	77% (13)					90% (39)	82% (38)	94% (53)	92% (25)	100% (8)	100% (2)	100%
PN-6	Initial antibiotic selection for CAP in immunocompetent patients	82% (11)	73% (11)					100% (15)	92% (13)	97% (36)	91% (23)	71% (7)	0 NA	100%

**Sedgwick County Medical Society
Core Measures Update:
1Q08 - 2Q08**

	Indicator	Galichia Medical Center		Kansas Heart Hospital		Kansas Spine Hospital		Via Christi		Wesley Medical Center		Kansas Medical Center		Ntl 90th %
		2Q08	3Q08	2Q08	3Q08	2Q08	3Q08	2Q08	3Q08	2Q08	3Q08	2Q08	3Q08	3Q07 - 2Q08
PN-6A	Initial antibiotic selection for CAP immunocompetent - ICU							100% (1)	50% (2)	100% (8)	71% (7)	0 NA	0 NA	NA
PN-6B	Initial antibiotic selection for CAP immunocompetent - Non ICU							100% (14)	100%	96% (28)	100% (16)	0 NA	0 NA	NA
PN-7	Influenza vaccination	NA	NA	NA	NA			NA	NA	NA	NA	NA	NA	97%
HF-1	Discharge instructions	97% (75)	98% (46)	96% (28)	94% (35)			62% (66)	62% (71)	80% (55)	76% (41)	78% (18)	84% (19)	98%
HF-2	Evaluation of LVS Function	93% (85)	98% (54)	100% (31)	100% (38)			97% (76)	100%	100% (70)	98% (49)	95% (22)	100% (21)	100%
HF-3	ACEI or ARB for LVSD	57% (35)	88% (17)	100% (8)	95% (19)			97% (35)	78% (27)	100% (16)	100% (15)	86% (7)	100% (9)	100%
HF-4	Adult smoking cessation advice/counseling	55% (11)	100% (8)	100% (4)	100% (2)			100% (18)	100%	100% (7)	100% (10)	100% (3)	100% (4)	100%
SCIP - Card 2	Surgery patients on beta blocker prior to admission who received beta blocker during periop period		97% (39)	89% (28)	92% (37)			91% (69)	91% (56)	95% (40)	100% (49)	79% (39)	86% (32)	NA
SCIP - 1A	Prophylactic antibiotic received within one hour prior to surgical incision -- overall rate	83% (59)	92% (66)	99% (68)	97% (67)			99% (148)	92% (149)	94% (114)	96% (147)	92% (79)	99% (76)	98%
SCIP - 2A	Prophylactic antibiotic selection for surgical patients - overall rate	100% (60)	100% (66)	100% (68)	100% (71)			99% (154)	100%	97% (117)	99% (150)	98% (80)	99% (77)	99%
SCIP - 3A	Prophylactic antibiotic discontinued within 24/48 hours after surgery end time - overall rate	88% (52)	97% (58)	89% (47)	83% (42)			94% (139)	89% (142)	91% (111)	93% (132)	70% (76)	74% (73)	98%
SCIP-VTE 1	Surgery patients with recommended VTE prophylaxis ordered	79% (24)	83% (12)	100% (3)	100% (3)			97% (123)	94% (123)	98% (132)	98% (163)	100% (45)	98% (53)	98%
SCIP - VTE 2	Surgery patients who received appropriate VTE prophylaxis within 24 hours prior to surgery to 24 hours after surgery	79% (24)	83% (12)	100% (3)	100% (3)			89% (123)	87% (123)	96% (132)	95% (163)	100% (45)	98% (53)	97%
SCIP - 4	Cardiac Surgery patients with controlled 6 A.m. post-op blood glucose		85% (66)	85% (52)	98% (56)					86% (35)	92% (39)	81% (47)	69% (42)	100%
SCIP - 6	Surgery Patients with Appropriate Hair Removal		92% (101)	96% (80)	100% (83)					95% (195)	91% (229)	99% (114)	99% (111)	100%

Value in parenthesis = "n" for indicator

AMI = Acute Myocardial Infarction

HF = Heart Failure

PN = Pneumonia

SCIP = Surgical Care Improvement Project